

Images of Interest / Imagens de Interesse

Pulmonary Cement Embolism*Embolia Pulmonar por Cimento Ósseo***Margarida Morgado^{1,2}, Ana Teresa Vilares^{1,2}, Filipe Barros Alves^{1,2}, Ricardo Castro¹, António J. Madureira^{1,2}**

¹Serviço de Radiologia, Centro Hospitalar Universitário de São João, Porto, Portugal.
²Departamento de Medicina, Faculdade de Medicina da Universidade do Porto, Porto, Portugal

Address

Margarida Morgado
Serviço de Radiologia
Centro Hospitalar Universitário de São João
Alameda Prof. Hernâni Monteiro
4200-319 Porto, Portugal
e-mail: margarida.r.morgado@gmail.com

Received: 04/09/2022**Accepted:** 05/10/2022**Published:** 30/04/2023

© Author(s) (or their employer(s)) and ARP
2023. Re-use permitted under CC BY-NC. No
commercial re-use.

Abstract

Vertebroplasty and kyphoplasty are widely used procedures for patients with symptomatic spinal compression fractures. Cement leakage into the paravertebral veins leading to pulmonary embolism is not an uncommon complication and the risk is increased when treating bone metastases. We present a case and images of a patient with previous vertebral compression fractures from metastatic breast cancer treated with vertebroplasty and kyphoplasty which presented imaging findings consistent with pulmonary cement embolism.

Keywords

Pulmonary embolism; Bone cements;
Vertebroplasty; Kyphoplasty; Fractures;
Compression.

Resumo

A vertebroplastia e a cifoplastia são procedimentos amplamente utilizados em doentes sintomáticos com fraturas vertebrais compressivas. O extravasamento de cimento ósseo para as veias paravertebrais com consequente embolia pulmonar não é uma complicação incomum e o risco é superior no tratamento de metástases ósseas. Apresentamos um caso clínico com imagens de uma doente com antecedentes de fraturas vertebrais compressivas por metastização óssea de cancro da mama, tratadas com vertebroplastia e cifoplastia, que evidenciou achados imagiológicos compatíveis com embolia pulmonar por cimento ósseo.

Palavras-chave

Embolia pulmonar; Cimentos ósseos;
Vertebroplastia; Cifoplastia; Fraturas;
Compressão.

Case

A 68-year-old female patient with previous vertebral compression fractures from metastatic breast cancer treated with vertebroplasty and kyphoplasty presented a new incidental thoracic finding on the surveillance computed tomography (CT) scan. Venous phase contrast-enhanced CT revealed high attenuating linear intravascular material within the left pulmonary artery and its posterior, descending and lateral basal branches (Fig. 1). The patient had no respiratory or cardiovascular symptoms.

Similar high attenuating linear intravascular material was found within the thoracic anterior external vertebral plexus at the level of T12 and the azygos vein (Fig. 2, A); and within the lumbar anterior external vertebral plexus at the level of L5, the left common iliac vein and the inferior vena cava (Fig. 2, B). The vertebral bodies of T11, T12 and L5 were filled with the same high-density material (Fig. 2).

These findings were consistent with pulmonary cement embolism after percutaneous vertebroplasty and kyphoplasty. (Fig. 3)

Discussion

Vertebroplasty and kyphoplasty are widely used minimally invasive procedures for patients with symptomatic spinal compression fractures.^{1,2,3} By injecting polymethyl methacrylate cement into the vertebral body, under image guidance, it provides vertebral stability and immediate




Figure 1 – Reconstructed oblique coronal plane venous phase contrast-enhanced CT (mediastinum window) with high attenuating linear intravascular material within the left pulmonary artery and its posterior and descending branches (dashed arrows) consistent with bone cement pulmonary emboli.

pain relief.^{1,3} Nevertheless, complications such as venous cement leakage into the paravertebral veins leading to pulmonary embolism are not uncommon.^{1,3} Compression fractures increase venous drainage of the already highly vascularized vertebral bodies³ and the risk of leakage is even higher when treating bone metastases since there is




Figure 2 – Reconstructed oblique sagittal plane venous phase contrast-enhanced CT (bone window) with high-density material filling the vertebral bodies of T11 and T12 and within the azygos vein (A, dashed arrow) and filling the vertebral body of L5 and within the left common iliac vein and the inferior vena cava (B, dashed arrow) indicating bone cement venous leakage.

often cortical destruction.¹ Furthermore, cement leakage into the inferior vena cava is a significant risk factor for pulmonary embolism.² Most patients with pulmonary cement embolism are asymptomatic and do not develop any sequelae.¹ Nevertheless, when associated with chest pain, dyspnoea, tachycardia or hypoxia, there should be suspicion of pulmonary infarction.^{2,3}

Conclusion

Pulmonary cement embolism is not an unusual complication associated with vertebroplasty and kyphoplasty. Radiologists

Ethical disclosures / Divulgações Éticas

Conflicts of interest: The authors have no conflicts of interest to declare.

Conflitos de interesse: Os autores declaram não possuir conflitos de interesse.

Financing Support: This work has not received any contribution, grant or scholarship.

Suporte financeiro: O presente trabalho não foi suportado por nenhum subsídio ou bolsa.

Confidentiality of data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Confidencialidade dos dados: Os autores declaram ter seguido os protocolos do seu centro de trabalho acerca da publicação dos dados de doentes.

Protection of human and animal subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Proteção de pessoas e animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos



Figure 3 – Postoperative thoracic and lumbar spine radiographs after vertebroplasty and kyphoplasty of T11, T12 and L5 demonstrate radio-dense linear opacities on the left peri-hilar region (arrow, A and B) and on the anterior paravertebral region (dashed arrow, B and C) consistent with leaked bone cement within the left pulmonary artery and its posterior and descending branches, the azygos vein, the left common iliac vein and the inferior vena cava.

should be able to recognize the characteristic intravascular and bone findings which allow a definite diagnosis, even when encountered incidentally.

responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial.

References

1. Choe DH, Marom EM, Ahrar K, Truong MT, Madewell JE. Pulmonary embolism of polymethyl methacrylate during percutaneous vertebroplasty and kyphoplasty. *AJR Am J Roentgenol.* 2004;183:1097-102.
2. Kim YJ, Lee JW, Park KW, Yeom JS, Jeong HS, Park JM, et al. Pulmonary cement embolism after percutaneous vertebroplasty in osteoporotic vertebral compression fractures: incidence, characteristics, and risk factors. *Radiology.* 2009;251:250-9.
3. Mansour A, Abdel-Razeq N, Abuali H, Makoseh M, Shaikh-Salem N, Abushalha K, et al. Cement pulmonary embolism as a complication of percutaneous vertebroplasty in cancer patients. *Cancer Imaging.* 2018;18:5.