

Elderly man with macroglossia and periorbital ecchymosis

Idoso com macroglossia e equimoses periorbitais

Mariana O. Fernandes^{1,a}, Luisa R. Ayoub^{1,b}, Samara O. Rabay^{1,c}, and Flávia R. Ferreira^{1,2,d}

¹Dermatologia, Hospital Municipal Universitário de Taubaté–H.MUT/UNITAU; ²Dermatologia, Departamento de Medicina da Universidade de Taubaté, UNITAU, Taubaté-SP, Brasil

ORCID: ^a0000-0002-3134-7096; ^b0000-0002-8461-1065; ^c0000-0003-1442-7040; ^d0000-0001-5679-4282

A Caucasian 70-year-old male was observed with macroglossia with hemorrhagic spots and dental impressions on the border of the tongue, eyelid ecchymosis (Fig. 1), multiple brownish, lichenified papules and plaques, and diffuse ecchymosis on the neck and upper trunk (Fig. 2). Lesions were pruritic and progressively developed during 6 months. No systemic symptoms, except fatigue.

Skin biopsy of a papular lesion on the upper trunk showed epidermal atrophy and diffuse dermal deposition of proteinaceous amorphous material colored with Hematoxylin & Eosin (HE) and further enhanced with Congo red stains (Fig. 3). This material showed green birefringence peculiar on polarized microscopy.

Protein electrophoresis showed a peak of gamma fraction and immunofixation showed a monoclonal IgG_κ with normal IgA and IgM. Bone marrow (BM) biopsy showed hypercellularity for age (in the three hematopoietic series) and the immunohistochemically panel showed BM infiltration by plasma cell neoplasia with evident lambda light chain restriction, confirming the diagnosis of multiple myeloma (MM). Neither osteolytic lesions nor the involvement of any other organ were detected.

The diagnosis of light chain (AL) amyloidosis associated with MM was done. The patient is being followed up together with onco-hematology staff, undergoing chemotherapy with Melphalan® and Prednisone, with

satisfactory evolution (important improvement in pruritus and mucosal lesions, stability of skin lesions, and the patient's general condition).

Discussion

Amyloidosis comprises a heterogeneous group of diseases with extracellular amyloid deposits¹. AL amyloidosis occurs in 5-26 % of patients with MM^{1,2}, and originates from the deposition in the skin and other organs of amyloid substance derived from monoclonal light chain immunoglobulins, leading to a plethora of clinical symptoms, and eventually vital organ dysfunction³. Lambda light chains are the most involved (3:1);³⁻⁵ however, in the present case kappa light chain was preponderant.

The age range of involvement is between 50 and 70 years, with a slight male predominance⁵. The clinical manifestations are variable and nonspecific (weight loss, fatigue, paresthesia, and syncope) contrasting with highly suggestive dermatological manifestations: macroglossia with tongue hardening, with or without bleeding spots and blisters in the oral cavity; periorbital ecchymosis (raccoon sign) and ecchymosis due to minor trauma in the folds; translucent, purpuric and/or brownish papules on the scalp, face, neck, and fingertips; which may be the only manifestations of the

Corresponding author:

*Flávia Regina Ferreira

Email: dermagica@uol.com.br

2797-5001 / © 2022 Portuguese Society of Dermatology and Venereology. Published by Permanyer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Received: 31-03-2022

Accepted: 27-05-2022

DOI: 10.24875/PJDV.M22000039

Available online: 24-10-2022

Port J Dermatol and Venereol. 2022;80(3):246-248

www.portuguesejournalofdermatology.com



Figure 1. A: ecchymosis on eyelids. B: macroglossia with dental impressions and bleeding spots.



Figure 2. Diffuse ecchymoses and brownish lichenified papules and plaques, in cervical region and trunk.

disease prior to later-stage organ involvement, at which point treatment options are limited¹⁻⁶.

All forms of amyloid show a peculiar green birefringence on microscopy after staining with Congo red, the gold standard for diagnosis^{5,6}.

In the present case, the histopathological examination (HE and Congo red) and the monoclonal peak of the gamma fraction on protein immunoelectrophoresis confirmed the diagnosis of AL amyloidosis. Hypercellularity for age in the BM biopsy and the

immunohistochemical panel showing BM infiltration by plasma cell neoplasm with evident lambda light chain restriction concluded AL amyloidosis was associated with MM.

As for the treatment, it aims to reduce the proliferation of the clonal B cell lineage and consequently the amyloid deposit, being the same used for MM (in the present case Melphalan® associated with Prednisone)^{1,6}. Prognosis reserved^{1,5,6}.

The rarity of this case (association), the importance of early diagnosis (minimizing/delaying the amyloid




Figure 3. Histopathological examination. **A and B:** positive protein material in HE. **C and D:** Red Congo stains.

deposit) and the fundamental role of the dermatologist in this context, motivated this report.

Acknowledgments

To Dr. Fátima Maria de Oliveira Rabay and the onco-hematology staff for the clinical management of this case and to Dr. Fernanda da Rocha Gonçalves for the diagnostic assistance and histopathological images.

Funding

None

Conflict of interests

None

Ethical disclosures

Protection of human and animal subjects. The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of

Ethics of the World Medical Association (Declaration of Helsinki).

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors have obtained the written informed consent of the patients or subjects mentioned in the article. The corresponding author is in possession of this document.

References

1. Al-Raddadi R, Abbaspour V, Ndiaye M. Papular cutaneous amyloidosis associated with multiple myeloma. *Science Journal of Clinical Medicine.* 2016;5:16–9. doi: 10.11648/j.sjcm.20160502.11
2. Hanumanthu V, Thakur V, Chatterjee D, Vinay K. Periocular cutaneous amyloidosis in multiple myeloma. *QJM.* 2021;114:603–4. doi: 10.1093/qjmed/hcab084
3. Mendelson L, Shelton A, Brauneis D, Sanchorawala V. AL amyloidosis in myeloma: red flag symptoms. *Clin Lymphoma Myeloma Leuk.* 2020;20:777–8. doi: 10.1016/j.clml.2020.05.023
4. Andrei M, Wang JC. Cutaneous light chain amyloidosis with multiple myeloma: a concise review. *Hematol Oncol Stem Cell Ther.* 2019;12:71–81. doi: 10.1016/j.hemonc.2018.09.003
5. Alcatrão MJ, Neves C, Gaspar A, Bravo A, Margarido E, Estrada H. Mieloma Múltiplo e Amiloidose AL. *Revista da Sociedade Portuguesa de Medicina Interna.* 2016;23:28–31.
6. Silverstein SR. Primary, systemic amyloidosis and the dermatologist where classic skin lesions may provide the clue for early diagnosis. *Dermatol Online J.* 2005;11:5.